2025

Exploring the social-cultural barriers of mental health among Hispanics

BOSTON UNIVERSITY

ARAM V. CHOBANIAN & EDWARD AVEDISIAN SCHOOL OF MEDICINE

Thesis

EXPLORING THE SOCIAL-CULTURAL BARRIERS OF MENTAL HEALTH AMONG HISPANICS

by

JONATHAN ROCHIN

B.A., California State University, Fresno, 2013

Submitted in partial fulfillment of the

requirements for the degree of

Master of Science

2025

Approved by

First Reader

Jonathan J. Wisco, Ph.D.
Associate Professor of Anatomy & Neurobiology

Second Reader

John Weinstein, Ph.D.
Director of Research

EPIGRAPH

"Every great dream begins with a dreamer. Always remember, you have within you the strength, the patience, and the passion to reach for the stars to change the world."

- Harriet Tubman

DEDICATION

I dedicate this work to my beloved spouse, Maria D. Rochin, for her unwavering support and for taking on the task of raising our wonderful children, Natalie and Isaac, while I am thousands of miles away. I also dedicate it to my parents, Teresa and David, who instilled in me the value of hard work and encouraged me to pursue my dreams against all odds.

ACKNOWLEDGMENTS

Thank you to the Boston University Aram V. Chobanian & Edward Avedisian Physician Assistant Program for giving me this opportunity. Special thanks Dr. Susan White, Dr. Jonathan Wisco and Dr. John Weinstein, for all their help and support.

EXPLORING THE SOCIAL-CULTURAL BARRIERS OF MENTAL HEALTH AMONG HISPANICS

JONATHAN ROCHIN

ABSTRACT

Mental health disparities among Latino populations in the United States remain a significant public health concern; Latinos experience lower access to mental health services and higher rates of treatment discontinuation compared to their white counterparts. Structural barriers, such as high costs, lack of insurance, and inadequate access to culturally competent care, contribute to these disparities. Cultural factors, including stigma surrounding mental health and differing perspectives on mental illness, further exacerbate the unmet needs within the community. For Latino immigrants in particular, socioeconomic challenges such as poverty and limited awareness of available services further restrict access to care. A critical gap exists between potential access to care and realized access with high rates of untreated mental illness among Latinos highlighting the urgency of addressing these issues.

Stigma, low mental health literacy, and cultural values such as familismo impede Latino individuals from seeking necessary mental health care. Interventions focused on enhancing social support and addressing unique coping mechanisms can improve mental health outcomes in this population. Culturally competent interventions, particularly those targeting healthcare providers, are essential to improve engagement and care for Latino patients. Prior research emphasizes the need for programs that integrate cultural

competence training, social support, and the recognition of cultural factors influencing health behaviors.

This study aims to develop and implement an educational program to enhance cultural sensitivity and competence among primary healthcare providers working with Hispanic patients. Grounded in Social Learning Theory and Transformative Learning Theory, the program will use a quasi-experimental pretest-posttest design to assess changes in healthcare providers' confidence and effectiveness in delivering culturally competent mental health care. Participants will be divided into an intervention group receiving 16 hours of training and a control group. Results will be compared to determine the impact of the training on providers' cultural competence.

Despite limitations such as potential selection bias and self-reporting, this study's design leverages proven adult learning theories to create a robust and practical intervention. By addressing cultural competence gaps, the study aims to reduce mental health disparities and improve the quality of care for Latino populations, ultimately contributing to greater health equity.

TABLE OF CONTENTS

LIST OF TABLES

LIST OF ABBREVIATIONS

CHIS .....California Health Interview Survey

CHW .....Community health workers

CME.....Continuous medical education

ED ..... Emergency Department

IBHC..... Integrated Behavioral Health Care

NAMI..... National Alliance on Mental Illness

PI.....Principal Investigator

RCT..... Randomized Controlled Trial

INTRODUCTION

Background

The Hispanic population in the United States is experiencing significant growth; it is the largest and fastest-growing minority group in the country.[1] Projections indicate that, by 2050, Hispanics will account for 19.8% of the older adult population.[1] This demographic shift highlights not only the vibrancy of Hispanic culture but also the unique challenges faced by this diverse group, including both U.S.-born and immigrant Hispanics, who encounter distinct obstacles related to their mental health care access.[1] As this population continues to expand, the mental health needs among Latinos in the United States become increasingly pressing. Research indicates that they experience mental health conditions at rates similar to or even higher than those of non-Latino whites.[2] However, despite this significant need, Latinos face substantial disparities in accessing and receiving adequate mental health care, marking a critical public health concern that demands urgent attention.[2] These disparities are rooted in both structural and cultural barriers that hinder their ability to obtain necessary services.[2] Addressing these barriers is essential not only for improving health outcomes within Latino communities but also for promoting overall equity in care delivery.

Statement of the Problem:

Structural barriers, including high costs of care, lack of health insurance, transportation difficulties, and a shortage of Spanish-speaking or culturally competent providers, create significant obstacles for Latino individuals seeking mental health services.[2] In addition to these structural impediments, cultural factors also play a crucial

role.[2] Stigma associated with mental health issues, differing cultural perspectives on mental illness, and fears related to immigration status often deter individuals from seeking help.[2]

These challenges are especially pronounced among older Hispanic adults, who experience a high prevalence of mental health issues such as depression, anxiety, and neurocognitive disorders.[1] These mental health conditions are often intertwined with physical disabilities, complicating treatment and exacerbating health challenges.[1] As such, the barriers to care not only impede access but also hinder the effectiveness of treatment for this vulnerable demographic.[1]

Moreover, recent sociopolitical developments, including changes in immigration policies, have further heightened the importance of examining mental health among immigrant populations.[3] The increasing stressors Latino families face, exacerbated by anti-immigrant rhetoric[3], can lead to heightened mental health challenges that necessitate urgent attention. Understanding and addressing these multifaceted issues is critical for improving mental health care access and outcomes for Latino communities.[3]

Hypothesis

Primary care healthcare providers who complete a minimum of 16 hours of social-competence training will demonstrate increased confidence and effectiveness in delivering mental health services to Hispanic patients compared to those who have not received such training.

Objectives and specific aims

This study aims to evaluate the impact of targeted interventions on enhancing healthcare providers' cultural competency with the goal of fostering greater socio-cultural awareness of the Hispanic community and identifying the barriers they encounter in accessing mental health services.

We will pursue the following two specific aims:

1. To identify and mitigate the social and cultural barriers within the Hispanic community that hinder access to mental health services.
2. To develop, implement, and evaluate a program that trains healthcare providers in a primary care setting in cultural sensitivity and competence

REVIEW OF LITERATURE

Overview

Mental health disparities among Latino populations in the United States remain a significant public health concern. Compared to their white counterparts, Latinos have lower rates of access to mental health services and are more likely to terminate treatment prematurely.[4] These disparities are intensified by structural barriers such as high costs of care, lack of health insurance, inadequate access to linguistically and culturally appropriate services, and transportation challenges.[4]

Additionally, cultural factors, including stigma surrounding mental health and differing perspectives on mental illness, contribute to significant unmet needs within the community.[4] Many individuals fear societal repercussions associated with seeking mental health support, which discourages them from pursuing necessary treatment.[5] Additionally, mental health literacy remains low in many Latino communities, leading to limited awareness of available resources and treatment options.[5] Cultural values such as *familismo,* which emphasizes strong family bonds, may further dissuade individuals from seeking outside help, as they may prefer to address challenges within the family unit rather than engage with external mental health services.[5]

For Latino immigrants, access to healthcare presents additional obstacles, as many face poverty, lack of health insurance, and limited awareness of available services.[6] It is crucial to distinguish between potential access (the availability of healthcare services) and realized access (actual use of these services) to effectively address the mental health

needs of this population.[6] Without bridging this gap, the high rates of untreated mental illness in Latino communities will continue to pose a pressing public health challenge.[6] Untreated mental health issues among Latinos can have significant consequences, affecting individuals and families alike.[5] Recognizing these challenges is essential for developing effective interventions aimed at improving mental health outcomes within this diverse population.

To make meaningful improvements in mental health outcomes among Latino immigrants, targeted interventions are necessary. These interventions should focus on enhancing social support systems and addressing the distinct mental health challenges faced by this community.[6] By understanding their unique coping mechanisms and social dynamics, healthcare providers can design services that align or resonate with the lived experiences of Latino individuals, increasing the likelihood of patient interaction.[6]

Given these complex issues, it is imperative that healthcare providers, particularly nurses and primary care professionals, take action to implement culturally competent mental health interventions tailored to the unique needs of Hispanic adults.[1] By fostering an inclusive and understanding healthcare environment, we can begin to bridge the gaps in mental health care access and ensure that this vulnerable population receives the comprehensive support necessary for their well-being. [1]

Existing Literature

Mental health remains a significant concern in the health sector. Despite ongoing efforts to address mental health challenges, there are still various factors hindering the

effectiveness in various communities. The Hispanic/Latino community have faced multiple barriers when accessing healthcare. These disparities have had a considerable impact on mental health intervention. Recognizing the importance of this issue, scholars have examined how culturally responsive practices can improve mental health access and outcomes for Hispanic/Latino populations.

Barriers and facilitators to depression care among Latino men in a primary care setting: A qualitative study

The study by Swetlitz et al.[4] highlights the unique and gender-related barriers Latino men face when accessing depression care. The study used semi-structured interviews with thirteen Latino men in primary care who were experiencing depressive symptoms. The participants were recruited through convenience and purposive sampling between December 2020 and August 2021. In identifying barriers and facilitators to depression care, researchers used consensual qualitative research and thematic analysis with modified grounded theory to analyze the data. This approach allowed for a culturally nuanced understanding of mental health barriers specific to Latino men. The authors' analysis identified various key barriers to depression care that relate to cultural perceptions of masculinity among Latino men. This included masculinity norms and pressure to be a provider, which discouraged the men from acknowledging depression or seeking help.

The qualitative design implemented in this study was key in allowing in-depth insights into cultural and structural barriers. The design, however, utilized a small sample

size, limiting generalizability. The subjective nature of qualitative data also presents some limitations to the approach, such as the introduction of potential bias based on individual experiences and researcher interpretation. These limitations suggest that the findings are more indicative than conclusive. However, the study offers valuable insights that could enhance cultural sensitivity training for providers by highlighting how cultural attitudes and masculinity norms impact Latino men's mental health care engagement. This approach may encourage clinicians to frame depression care as congruent with positive masculine values.

Social support, health, and health care access among Latino immigrant men in an emerging community. Health Education & Behavior

Further examining social support, the study by Document et al.[6] highlighted how social support is significantly associated with reduced binge drinking and lower rates of depression, despite not being readily available. The study employed a cross-sectional design that gathered data between 2011 and 2013. A notable strength of the study was the process involved recruiting eleven male Latino community health workers (CHWs) from various community settings to deploy a 90-question baseline survey to 682 participants (Latino immigrant men), of which a sample size of 125 was used for the evaluation.[6] This approach likely enhanced the honesty of participant responses, as they were more comfortable interacting with CHWs who shared their gender and cultural background. Logistic regression analyses were performed to investigate the relationships between social support and health behaviors. The analysis revealed that Latino immigrant men in

the study were generally at high risk of mental health issues based on their low educational attainment.[6] Their risks entail high rates of depression and limited access to healthcare (only 6% had health insurance).[6] Higher social support within this population was associated with 78% fewer cases of moderate-to-severe depression and 56% fewer cases of binge drinking.

This study highlights important findings on how social determinants like social support can influence health behavior among vulnerable communities. Focusing on Latino men allowed the study to fill the gaps in understanding how underserved immigrant groups cope with stress and limited resources. However, the study's cross-sectional design makes it difficult to prove causation because the data were gathered all at once without a longitudinal follow-up to track changes in health outcomes. Another strength of the study is the community-based participatory approach that involves the locals as CHWs. This approach ensured relevance and impact on the findings by addressing the unique challenges faced by Latino immigrants. While the reliance on self-reported biases – particularly on sensitive topics like drinking behavior and depression – the researchers' decision to employ male Latino CHWs to perform the surveys likely minimized these biases as much as possible. Another limitation of the study is the restriction of the sample to Latino immigrant men.[6] This limitation hinders the generalizability of the findings to Latino populations in other regions.

Health care access and physical and behavioral health among undocumented Latinos in California. Medical Care

The need for culturally and socially informed healthcare approaches is reinforced by Ortega et al.[7] whose study highlights the role of social and cultural competence in delivering mental health interventions to Latinos. Their study employed data from the California Health Interview Survey (CHIS) covering the period between 2011 and 2015. The researchers investigated the comparative aspects of healthcare access, utilization and health status of nonelderly Latino and U.S.-born non-Latino white adults in California. In the analysis, participants were grouped based on citizenship status, allowing the use of multivariable regression models to examine associations between citizenship/documentation status and health access, utilization, and physical and behavioral health outcomes among the groups. The research identified four large variations based on documentation of the status of health care being available, used, and affordable to nonelderly Latino and U.S.-born non-Latino white adults in California.[7] The groups included U.S-born Latinos: 8,303 participants, Naturalized Latinos: 3,879 participants, Green Card Holders: 3,369 participants, and Undocumented Latinos: 3,053 participants.[7] Based on the data indicating basic connections between citizenship or documentation status and physical and mental health, unmet health needs, and access to healthcare, Latino men have the lowest healthcare use and face the most significant barriers to accessing healthcare compared to other groups.[7] Only 61% of the participants had a regular source of healthcare outside of the Emergency Department (ED), meaning may relied on the ED for healthcare needs or had no regular care at all.[7] They also were the least likely to visit the ED (14%) and the least likely to have seen a doctor (58%),

reporting the fewest doctor visits over the year, indicating limited overall interaction with healthcare services.[7]

These findings highlight the disparities that exist in healthcare access and utilization among undocumented Latinos. The authors suggest that, to have an effective intervention, healthcare professionals should first focus on understanding the cultural and social status of the community, including the barriers they face. This competency helps ensure that mental health interventions are inaccessible due to systemic barriers among the target population. The strengths of this study design include a large, representative sample and detailed stratification of documentation status. This advantage helped in providing valuable insights across various citizenship categories. However, the study's reliance on self-reported data appeared as a limitation, which could lead to underreporting of health conditions, especially among undocumented participants, due to fear or stigma. Additionally, the focus on California may limit the study's applicability to Latino populations in other states. In states with stricter immigration policies and fewer safety-net programs, undocumented Latinos likely face even greater barriers to healthcare.[7] They may experience lower access and utilization rates compared to California, where supportive county programs slightly improve healthcare accessibility.[7] Overall, this study points out the need for policies that improve healthcare access for undocumented Latinos, facing significant barriers.

A parish-based multilevel cluster randomized controlled trial to reduce stigma and mental health treatment disparities among Latino communities. Contemporary Clinical Trials

Building on these themes of cultural competency, Wong et al.[5] note that cultural factors, low mental health knowledge, and stigma contribute to the sizable percentage of the Latino population in the United States going undiagnosed and untreated. Therefore, in the article, the Diocese of San Bernardino and the National Alliance on Mental Illness (NAMI) attempt to address the issue of inequalities among Latino parishioners through use of religious congregations. The design for this study is a cluster Randomized Controlled Trial (RCT). It matched and assigned parishes based on size and language preferences into two groups (intervention or control). Implementation will be phased over two waves, with surveys conducted at baseline, six, and twelve months to evaluate results. Participants will be recruited from parishes during religious services and will complete surveys in either paper or web formats. The primary outcomes of this study focus on mental health service usage and support for those with mental health issues. Additional measures include mental health knowledge, perceived treatment needs, stigma, and structural barriers, using validated tools like the Mental Health Literacy Scale and Perceived Need for Care Questionnaire.

Wong et al.[5] article supports the hypothesis and the objectives by highlighting the importance of cultural competency in addressing barriers faced by Hispanic patients, thus validating the need for targeted interventions to enhance social-cultural awareness and accessibility of mental health services. The RCT design used is a key strength of the

study. It minimizes bias and the risk of confounding and allows for clear comparisons between intervention and control groups. This strengthens the validity of conclusions about the effectiveness of social-competence training. However, the design may have limitations in generalizability. As the study focuses on a single geographic area-California's Riverside and San Bernardino Countries, it might overlook other areas with high Latino populations.[5] The findings in the study might not fully apply to them based on different socioeconomic or health service dynamics

Coping with mental health issues among older Hispanic adults

Adding further depth to these findings, Curtin et al.[1] examine coping strategies used by elderly Hispanic immigrants to manage mental health concerns, such as stress, nervousness, and depression. The study used a cross-sectional, descriptive research design, where 17 older Hispanic adults who attended a low-income clinic were interviewed. Participants included those who were 65 years and above and had migrated and lived in the area for four to thirty-nine years. Semi-structured interviews were also employed to determine the influence of beliefs and experiences on the coping mechanisms.

The findings revealed that older Hispanic immigrants cope with mental health issues through professional assistance, social support, spiritual beliefs, and diversion. Many of the participants relied on their faith in God, viewed social connections with family and friends as crucial, and employed various distraction techniques to manage stress, anxiety, and depression. These results demonstrate that older immigrants employ

culturally relevant coping mechanisms, such as spirituality and social support, to manage mental health challenges. These strategies align with the objectives of understanding how cultural factors influence coping behaviors. Therefore, cultural competencies are crucial in addressing mental health issues among Hispanics.

This study design has various strengths, such as the focus on a specific cultural group. This approach provides unique coping mechanisms that are crucial in noting the conclusions. The qualitative methods also allow for an in-depth exploration of participant opinions, thus validating the results provided. The study, however, has some limitations, such as a small sample size and potential selection bias, which may affect the findings' representativeness. While the insights are valuable for understanding this particular demographic, it is crucial for further research to involve a much more diverse sample to enhance generalizability across different immigrant populations and mental health contexts.

Does integrated behavioral health care reduce mental health disparities for Latinos? Initial findings

In a related investigation, Bridges et al.[2] evaluated the effectiveness of Integrated Behavioral Health Care (IBHC) for Latino patients - a model designed to lessen stigma and obstacles related to mental health services by incorporating mental health professionals within primary care—on mental health differences between patients who are Latino and those who are not. It analyzed the referral, utilization, and outcome patterns of 793 patients (63.8% Latino) who sought behavioral health services in two

medical clinics over 10 months. The clinics operate on a sliding fee scale and provide care irrespective of patients' insurance status. Information was gathered from patients who received mental health services in succession from August 2011 to June 2012. Out of 1,964 behavioral health patients, 823 had complete data. The study found that, compared to non-Latino White patients, Latino patients showed higher levels of self-reported psychological distress. However, Latinos were less likely to receive a psychiatric diagnosis.[2] These findings align with the hypothesis that culturally competent professionals can craft interventions that override the barriers affecting the Latino population.

The study's design strengths include its focus on culturally tailored interventions through Integrated Behavioral Health Care. This approach enhances relevance for Latino populations and addresses specific mental health needs. However, the design also has weaknesses, such as a limited sample size. The small sample size affects generalizability and possible selection bias. This is because the participants may already be inclined toward treatment. The lack of diversity within the Latino sample might also limit broader applicability across varied Latino subgroups[2], because the study primarily focused on Spanish-speaking Latino patients, which may not represent the broader Latino population. Additionally, cultural variations in how distress is expressed across Latino subgroups were not fully explored, limiting the generalizability of the findings.[2] In addressing these limitations, further studies can refine the approach by enhancing diversity and sample size, thus expanding our understanding of culturally competent care's impact.

Conclusion

In summary, the findings across these studies collectively highlight the importance of culturally competent interventions in mental health care for Latino populations. The literature reveals common barriers to mental health access for Hispanic and Latino populations (Table 1), by highlighting themes such as limited social support, high stigma, and cultural norms. These factors intersect to create significant obstacles in recognizing, seeking and receiving appropriate medical care. The validity of these results is highlighted by the diverse research design employed in the studies.

Despite the limitations, such as small sample sizes, generalizability, and subjective data emerging from the study design, cultural sensitivity emerges as a crucial factor in improving access to and outcomes of mental health services for these populations. This highlights the need for further research to build on and expand these findings. The implication of these findings stresses the urgent need for culturally tailored interventions to improve mental health outcomes, foster socio-cultural awareness, and address specific barriers faced by Hispanic patients.

Table 1: Common Barriers to Mental Health Access for Latino Populations

Study	Populations	Key Barriers Identified	Support Systems / Facilitators
Article: Depression Care Among Latino Men	Latino men in primary care settings	Masculinity-related barriers (emotional control, self-reliance), stigma around help-seeking	Encouraging honesty, collaboration, responsibility in care
Article: Social Support, Health, and Health Care Access	Encouraging honesty, collaboration, responsibility in care	Low health care access, low insurance, high rates of depression and binge drinking	Social support reduces depression and binge drinking
Article: Health Care Access and Physical and Behavioral Health	Undocumented Latinos in California	Limited access to healthcare, legal and financial barrier	Social and community networks
Article: Parish-Based Intervention to Reduce Stigma	Latino religious communities	Mental health stigma, lack of mental health literacy, reluctance to seek help outside religious settings	Religious congregations, community partnerships
Article: Coping Mechanisms in Older Hispanic Adults	Older Hispanic immigrants from Guatemala, Dominican Republic, Colombia	Limited access to mental health services, reliance on self-management	Spirituality, social support, primary care providers
Article: Integrated Behavioral Health Care (IBHC)	Latino and non-Latino White primary care patients	Stigma, barriers to mental health service utilization	Integration of behavioral health into primary care

METHODS

STUDY DESIGN

The goal of this study is to develop and implement an educational program aimed at enhancing cultural sensitivity and competence among primary healthcare providers. This learning initiative is grounded in Social Learning Theory and Transformative Learning Theory. Social Learning Theory, proposed by Albert Bandura, highlights that individuals acquire knowledge through observation, imitation, and modeling.[8] According to this theory, primary healthcare providers can build cultural sensitivity by observing culturally competent behaviors, participating in role-playing scenarios, and receiving feedback from peers and mentors.[8] Transformative Learning Theory, developed by Jack Mezirow, focuses on adult learning, emphasizing that individuals can revise their perspectives based on new insights.[8] This theory proposes that healthcare providers can transform their approaches by critically reflecting on their experiences, assumptions, and biases, leading to a deeper understanding and practice of cultural competence.[8]

To achieve the study's objectives, researchers will employ a non-random allocation method, consistent with the quasi-experimental pretest-posttest design to assess the impact of the social-competence training program on providers' confidence and effectiveness in working with Hispanic patients. Participants will include primary healthcare providers, divided into an intervention group receiving 16 hours of social-competence training and a control group that does not participate in the training. The intervention group will complete feedback forms after each training module, allowing researchers to collect qualitative data on their experiences and suggestions for

improvement. Both groups will complete a pretest assessment (Appendix A) to establish baseline measures of confidence and effectiveness in mental health service delivery for Hispanic patients. After the intervention, a posttest assessment (Appendix B) will be conducted for both groups using identical metrics, allowing researchers to evaluate changes and compare outcomes within and between the groups.

Study Population and Sampling

The source population for this study consists of primary healthcare providers who work with Hispanic patients in Los Angeles, California. These providers include professionals who are directly involved in patient care, such as physicians, physician assistants, nurse practitioners, and nurses, particularly those serving in primary care or community health settings where a significant portion of the patient population identifies as Hispanic.

Participants will be sampled from this source population using a convenience sampling method. Invitations to participate in the study will be sent to primary healthcare facilities and community health centers known for their engagement with Hispanic patients. Providers who meet the inclusion criteria will be invited to participate until the desired sample size is reached.

The inclusion criteria for this study specify that participants must be primary healthcare providers, such as physicians, physician assistants, nurse practitioners, and nurses, who actively work with Hispanic patients in Los Angeles, California. Additionally, eligible providers should have at least six months of experience in a

primary or community healthcare setting to ensure familiarity with patient care routines. Participants must also be willing to engage for the full study duration, including completing both pretest and posttest assessments, as well as attending intervention sessions if assigned to the training group.

The sample size for the study was calculated as 199 per arm using the *Difference between Two Means (Paired Samples T-Test) calculator* from the Statistical Power and Sample Size Calculation Tools.[9] The calculation was based on Cohen's d=0.2, a correlation between repeated measures of 0.5, a statistical power of 0.8, an alpha level of 0.05, and an alternative hypothesis of "Not Equal." Using a small effect size (Cohen's d=0.2) is a conservative and appropriate choice in this context, as there is no prior evidence to predict how effective the intervention will be. By assuming a small effect size, the study is powered to detect a wide range of effect sizes while maintaining a feasible sample size. This approach ensures that even subtle improvements can be identified without requiring an impractically large participant pool. This sample size ensures a 95% confidence level, minimizing the risk of Type I errors.[9] The primary outcome measure is the change in healthcare providers' self-reported confidence and effectiveness in working with Hispanic patients, evaluated using a standardized cultural competence assessment scale. This tool quantifies improvement by comparing pretest and posttest results within and between the intervention and control groups.

Intervention

The intervention for this study, a social-competence training program for healthcare providers, is in the developmental stage and will be designed specifically to enhance providers' skills in culturally responsive care for Hispanic patients. It will aim to build cultural awareness, improve communication techniques, and address key barriers to care that Hispanic patients often face. To maximize accessibility and flexibility, the training sessions will be conducted digitally as live events. These sessions will also be recorded, allowing participants who are unable to attend the live training to review the material at their convenience. This approach ensures that all participants have the opportunity to engage fully with the program, regardless of scheduling constraints.

Key elements proposed for the program include modules on *Cultural Awareness and Sensitivity*, where providers will receive information on the diversity within Hispanic communities and learn about core cultural values like *familismo*, *respeto*, and *personalismo*, which can significantly influence patient interactions and health decisions. In the *Effective Communication Skills* module, providers will practice techniques like using interpreters, employing active listening, and understanding nonverbal communication differences to build rapport and trust with Spanish-speaking patients.

The program will also address *Barriers to Care* by training providers to recognize and mitigate challenges Hispanic patients may face, such as limited access to transportation or financial resources, and by offering strategies to connect patients to community support. Additionally, the module on *Understanding Mental Health in Context* will cover cultural perspectives on mental health, tackling common

misconceptions, and educating providers on the prevalence of mental health conditions like depression, anxiety, and neurocognitive disorders.[1]

To reinforce practical skills, the training will incorporate *Scenario-Based Learning* through role-playing exercises and case studies, allowing providers to apply culturally competent practices in simulated patient interactions. By integrating these elements, the training program aims to empower healthcare providers to deliver more effective, culturally tailored care, which in turn can lead to better health outcomes and reduce healthcare disparities for Hispanic communities.

Study Variable and Measures

In this study, the primary variables include the independent variable—participation in the social-competence training program—and the dependent variables, which are the healthcare providers' self-reported *confidence* and *effectiveness* in delivering culturally competent care to Hispanic patients.

The independent variable, *social-competence training*, will be operationalized by providing the intervention group with a structured 16-hour training program. This program will include modules on cultural awareness, effective communication skills, barriers to care, mental health perspectives, and scenario-based learning, specifically tailored for working with Hispanic patients. The control group will not receive this training, allowing researchers to compare outcomes between those who did and did not receive the intervention.

The dependent variables, *confidence* and *effectiveness*, will be measured using a standardized cultural competence assessment tool designed to capture providers' self-assessed skills and attitudes. This tool will be administered as both a pretest and a posttest to measure any changes attributable to the intervention. Confidence refers to the providers' self-perceived ability to interact effectively and sensitively with Hispanic patients, while effectiveness pertains to their self-assessed capability to provide culturally competent care that aligns with the needs and values of this population.

The study will use an alpha level of 0.05 to determine statistical significance and assess meaningful changes in confidence and effectiveness scores from pretest to posttest within and between the intervention and control groups. By analyzing these variables, the study aims to determine the impact of social-competence training on healthcare providers' abilities to deliver culturally responsive care, thereby addressing healthcare disparities and improving outcomes for Hispanic communities.

Recruitment

To effectively recruit participants for this study, we will target primary healthcare providers working with Hispanic patients in Los Angeles, California.

Recruitment efforts will focus on primary healthcare facilities and community health centers renowned for their engagement with the Hispanic community. Initially, we will identify these institutions through local healthcare directories, professional networks, and partnerships with healthcare organizations that serve a significant Hispanic

population. Once identified, invitations to participate will be sent via email and postal mail to healthcare administrators and directly to providers, outlining the study's objectives, benefits, and requirements.

Additionally, informational flyers and brochures will be distributed within these facilities to reach potential participants directly. To enhance our engagement, we will collaborate with facility leaders to present brief informational sessions during staff meetings, highlighting the importance of cultural competence and the study's potential impact on patient care. Providers who meet the inclusion criteria—such as having at least six months of experience in a primary or community healthcare setting and actively working with Hispanic patients—will be invited to enroll in the study.

To encourage participation and ensure commitment, we may offer incentives such as continuing medical education (CME) credits or small monetary rewards upon completion of the study. Throughout the recruitment process, clear communication will be maintained to address any questions or concerns, ensuring that participants are fully informed and willingly consent to participate.

Data Collection

Data for this study will be collected through pretest and posttest assessments completed by participants in both the intervention and control groups. These assessments will be deployed via email using REDCap, a secure, HIPAA-compliant online survey platform. Each assessment will include a standardized cultural competence scale

designed to measure participants' self-reported confidence and effectiveness in delivering culturally competent care to Hispanic patients.

Additionally, the intervention group will complete feedback forms after each training module, providing qualitative data on their experiences and perceived benefits of the training program. To ensure secure and anonymous data collection, each participant will be assigned a unique identifier. This identifier will link their pretest and posttest results without requiring identifiable information. For qualitative data collected through feedback forms, responses will also be linked to these unique identifiers while preserving participant anonymity. Electronic assessments will allow participants to complete surveys conveniently on personal or work devices, ensuring flexibility and accessibility throughout the data collection process.

Once collected, all electronic data will be stored on a secure, password-protected server with access restricted to authorized research personnel. The data will be encrypted to safeguard against unauthorized access, and backup copies will be maintained in a separate, secure location to prevent data loss. To further protect participant confidentiality, any hard copies of data, if generated, will be stored in a locked cabinet within a secure research facility, and digital files will be de-identified before analysis.

Data retention will comply with institutional and federal guidelines, with all records maintained for five years post-study to allow for potential follow-up analysis or review. After this period, electronic data will be permanently deleted, and any hard copies will be shredded to ensure participant confidentiality.

Data analysis

The data for this study will be reviewed and analyzed using a combination of descriptive, inferential, and qualitative analysis methods to comprehensively assess the impact of the social-competence training program. Descriptive statistics, including the mean and standard deviation of each pre-training and post-training assessment question, will provide a baseline understanding of average scores and response variability. These descriptive metrics will highlight overall trends and changes in participants' knowledge and confidence levels.

For inferential analysis, a paired t-test will be conducted on pre-test and post-test scores if the data is normally distributed. This test will help determine whether there are statistically significant differences in scores after training, indicating a potential positive effect of the intervention. If the data is not normally distributed, the Wilcoxon signed-rank test, a non-parametric alternative, will be used. Additionally, effect size, calculated using Cohen's d, will provide context for the magnitude of change, offering insights into the practical significance of any observed differences.

For open-ended feedback, thematic analysis will be applied. This will involve systematically reviewing responses, coding key themes, and grouping similar codes into broader categories, ultimately capturing participants' reflections on the training's relevance, areas for improvement, and overall experience. This qualitative analysis will offer a deeper understanding of how participants perceive the program and provide specific feedback for refining the intervention.

The results will be summarized in a comprehensive report, including statistical outcomes (e.g., p-values, means, effect sizes) and insights from thematic analysis. Based on this review, recommendations for future training modifications or enhancements will be offered to improve the program's effectiveness. Stratified analysis, if needed, will consider subgroups, such as years of experience or specific roles within healthcare, allowing for targeted insights and identifying how the training may impact different provider demographics.

Timeline and Resources

The project to enhance cultural competence among healthcare providers in Los Angeles will unfold over an estimated 12-month period. During the initial three months, the focus will be on finalizing the design and content of the social-competence training program, which includes modules on Cultural Awareness, Effective Communication, Barriers to Care, and Understanding Mental Health in Context. During this stage, materials such as role-playing scenarios, case studies, and instructional guides will be developed. In addition, validated assessment tools and feedback forms will be selected to measure program effectiveness.

Months four and five will center on participant recruitment and pretest assessment. Invitations will be distributed to targeted healthcare facilities, and informational sessions will be held to inform potential participants about the study's goals and the importance of cultural competence. Eligible providers who consent will

complete a pretest assessment to establish baseline measures of confidence and effectiveness in culturally responsive care.

Implementation of the intervention will take place in months six and seven, during which the intervention group will participate in a 16-hour training program conducted over multiple sessions. Each session will feature various training activities, including role-playing and case studies, followed by feedback forms to collect qualitative data from participants regarding their experiences with each module.

In month eight, both the intervention and control groups will complete a posttest assessment to gauge any changes in confidence and effectiveness. During this phase, all data will be reviewed for accuracy and completeness.

Data analysis and interpretation will occur over a three-month period during months nine, ten, and eleven. Quantitative data will be evaluated through descriptive and inferential statistics to identify any statistically significant improvements. Qualitative feedback will be examined using thematic analysis to provide in-depth insights into participants' perceptions and suggestions for program enhancement. Visual presentations of the results, such as charts and word clouds, will be prepared to aid in communicating findings.

The final month, month twelve, will be dedicated to reporting and disseminating the results. The project findings will be summarized in a comprehensive report, which will include recommendations for program improvement and scalability, and presented to stakeholders.

The project requires key personnel to ensure its success. A Principal Investigator (PI) will oversee the project, manage compliance, and coordinate the team. A Project Coordinator will handle daily tasks such as scheduling, participant communication, and ensuring data is collected accurately. A Data Analyst will manage quantitative analysis, while a Qualitative Research Analyst will handle thematic analysis of feedback. Training Facilitators (two to three individuals) will lead the sessions, guide role-playing exercises, and engage participants. Additionally, Technical Support Staff will ensure smooth operation of the online survey platform and data security. Finally, an External Consultant with expertise in cultural competence will assist with program content development, particularly regarding Hispanic cultural values and healthcare disparities.

Institutional Review Board

Based on the BUMC IRB website, the study falls under Exempt Category 2.2 and will require the IRB to conduct a "limited IRB review."[10] While the study involves healthcare providers discussing potentially sensitive topics and addresses mental health outcomes for Hispanic populations, participation will be entirely voluntary, with participants providing informed consent. All data will be handled with strict confidentiality to protect participant privacy.

CONCLUSION

Discussion

This study has several limitations that could affect its findings and generalizability. First, the convenience sampling method used to recruit participants may introduce selection bias, as only providers from certain facilities will participate. This could impact the representativeness of the sample, limiting the generalizability of results to other regions or healthcare settings. Additionally, self-reported measures of confidence and effectiveness in cultural competence may be subject to social desirability bias, as providers might overstate their cultural competence or effectiveness in working with Hispanic patients. This reliance on subjective assessments could potentially skew results, making it challenging to capture an accurate measure of the intervention's impact.

Moreover, the non-random allocation method, consistent with the quasi-experimental design with a control group that does not receive the intervention but is not randomly assigned may introduce internal validity concerns. Differences between groups other than the intervention could influence outcomes, potentially confounding the findings. To address this, baseline assessments are conducted, but the lack of randomization means that group differences cannot be entirely ruled out.

Despite these limitations, the study also has notable strengths. A key advantage is its grounding in Social Learning Theory and Transformative Learning Theory, which are well-suited to adult learners in professional settings. By incorporating elements such as observation, role-playing, and reflective practice, the training program aligns with proven adult learning principles, potentially enhancing its effectiveness in promoting sustained

behavioral change. Furthermore, the 16-hour training program is comprehensive, addressing multiple facets of cultural competence, from communication skills to understanding mental health in a cultural context. This multi-dimensional approach is likely more robust than existing studies that focus on a single aspect of cultural competence.

An anticipated obstacle is the time commitment required for the intervention; busy healthcare providers may find it challenging to commit to 16 hours of training. To mitigate this, the program could be offered as continuing medical education (CME) credits to incentivize participation. Additionally, coordinating logistics for data collection, especially ensuring pretest and posttest completion from all participants, may be challenging in a clinical environment with high demands on providers' time.

Ultimately, this study offers a structured, theory-based training model that not only aims to increase cultural competence among providers but also attempts to address healthcare disparities within the Hispanic community. The findings, while limited in their generalizability, have the potential to contribute valuable insights into effective training approaches for culturally responsive care. By focusing on primary healthcare providers in Los Angeles—a region with a significant Hispanic population—this study addresses a critical need, and its results may inform future interventions aimed at reducing disparities and improving healthcare quality for minority populations.

Summary

The literature review underscores the persistent mental health disparities faced by Latino populations in the United States. Key findings indicate that Latino individuals, particularly men and immigrants, face unique structural and cultural barriers, such as high care costs, lack of health insurance, and stigma associated with mental health. These factors are exacerbated by limited access to culturally competent care and compounded by cultural norms like *familismo* and masculinity pressures that deter individuals from seeking help outside their immediate family. Stigma and low mental health literacy are critical challenges that further impede timely care, highlighting the need for targeted mental health education within the community.

Research also reveals that social support systems, when available, are linked to better mental health outcomes, such as reduced depression and binge drinking rates. However, accessibility remains a challenge for immigrant Latino men, who often lack adequate social resources and health insurance. Additionally, specific coping strategies, including reliance on spirituality and social bonds, demonstrate the cultural uniqueness of Latino populations in managing mental health, indicating that cultural competence in healthcare is paramount. Studies show that understanding Latino values, like family cohesion and faith, can help healthcare providers offer more relevant and effective mental health interventions.

The use of diverse methodologies across the studies provides valuable insights into these challenges. Qualitative approaches allow for an in-depth understanding of specific barriers, such as cultural perceptions of masculinity, while quantitative studies highlight broader patterns, such as the association between social support and health

outcomes. Despite limitations, including sample size constraints, generalizability issues, and the reliance on self-reported data, the studies emphasize the importance of culturally tailored care in overcoming the unique mental health barriers experienced by Latino populations.

In conclusion, there is an urgent need to implement culturally sensitive mental health interventions that acknowledge and address the complex social and cultural factors affecting Latino communities. Targeted efforts by healthcare providers, particularly those in primary and community health settings, could enhance engagement and reduce stigma. By fostering an inclusive, culturally aware healthcare environment, interventions can effectively bridge the existing gaps in mental health care access and outcomes, leading to improved well-being for Latino populations.

Clinical and/or Public Health Significance

Hispanics in the United States face substantial barriers in accessing and utilizing mental health services, resulting in higher rates of untreated mental illness compared to other demographic groups. These barriers include cultural beliefs, language obstacles, stigma, and structural inequalities such as lack of insurance and financial constraints. Culturally competent care is critical in mitigating these disparities. Studies reveal that Hispanics, especially those with a strong ethnic identity and limited English proficiency, are less likely to seek mental health services. This reluctance is often influenced by

cultural values that emphasize family cohesion and collectivism, which can discourage individuals from seeking help due to concerns about family reputation.

Language barriers exacerbate the issue, as the shortage of bilingual mental health professionals hinders effective communication and treatment. Stigma surrounding mental illness in Hispanic culture further discourages individuals from seeking care, contributing to the underutilization of mental health services.

These disparities highlight the need for targeted interventions that address the unique cultural and linguistic needs of the Hispanic community. By promoting culturally competent care and reducing structural barriers, healthcare systems can improve mental health outcomes and ensure equitable access to mental health services for Hispanics. Addressing these challenges is essential for reducing the public health burden of untreated mental illness in this population and promoting overall health equity.

APPENDIX

APPENDIX A: **Pre-Test Assessment**

Section 1: Knowledge Assessment

1. **Multiple Choice Questions:**

    1. What percentage of the Hispanic population in the U.S. is foreign-born?

        - A) 25%
        - B) 37%
        - C) 50%
        - D) 60%

    2. Which of the following cultural values is commonly associated with the Hispanic community?

        - A) Individualism
        - B) Familismo
        - C) Stoicism
        - D) Secularism

    3. Which mental health disorder is most commonly reported among older Hispanic adults?

        - A) Schizophrenia
        - B) Bipolar Disorder
        - C) Depression
        - D) Eating Disorders

2. **True/False Questions:**

    1. Hispanic individuals are less likely to seek mental health treatment due to stigma associated with mental illness. (True/False)

    2. Language barriers have no impact on access to healthcare for Hispanic patients. (True/False)

Section 2: Confidence Assessment

On a scale of 1 to 5, rate your confidence in the following areas (1 = not confident, 5 = very confident):

1. Understanding cultural values and beliefs of Hispanic patients.
2. Communicating effectively with Hispanic patients who may have limited English proficiency.
3. Identifying mental health issues in Hispanic patients.
4. Providing culturally sensitive care to Hispanic patients.

APPENDIX B: **Post-Test Assessment**

Section 1: Knowledge Assessment

1. **Multiple Choice Questions:**

    1. What is an important cultural value in Hispanic communities that impacts healthcare?

        - A) Autonomy
        - B) Familismo
        - C) Individualism
        - D) Aggression

    2. What is a common barrier to accessing mental health services for Hispanic individuals?

        - A) Lack of interest
        - B) High cost of services
        - C) Preference for alternative medicine
        - D) All of the above

    3. How can primary care providers best support Hispanic patients seeking mental health services?

        - A) Providing referrals only
        - B) Involving family in the treatment process
        - C) Offering only medication management
        - D) None of the above

2. **True/False Questions:**

    1. The majority of Hispanic patients prefer to communicate with their healthcare provider in English. (True/False)

    2. Understanding the cultural context of mental health can improve treatment adherence among Hispanic patients. (True/False)

Section 2: Confidence Assessment

On a scale of 1 to 5, rate your confidence in the following areas (1 = not confident, 5 = very confident):

1. Understanding cultural values and beliefs of Hispanic patients.
2. Communicating effectively with Hispanic patients who may have limited English proficiency.
3. Identifying mental health issues in Hispanic patients.
4. Providing culturally sensitive care to Hispanic patients.

BIBLIOGRAPHY

1. Curtin A, Martins DC, Schwartz-Barcott D. *Coping with mental health issues among older Hispanic adults.* Geriatric Nursing (Lond). 2019;40(2):123-128. doi:10.1016/j.gerinurse.2018.07.003

2. Bridges AJ, Arthur R Andrews III, Villalobos BT, Pastrana FA, Cavell TA, Gomez D. *Does Integrated Behavioral Health Care Reduce Mental Health Disparities for Latinos? Initial Findings.* Journal of Latina/o Psychology. 2014;2(1):37. doi:10.1037/lat0000009

3. Bucay-Harari L, Page KR, Krawczyk N, Robles YP, Castillo-Salgado C. *Mental Health Needs of an Emerging Latino Community.* Journal of Behavioral Health Services & Research. 2020;47(3):388. doi:10.1007/s11414-020-09688-3

4. Swetlitz N, Hinton L, Rivera M, Liu M, Fernandez AC, Garcia ME. *Barriers and facilitators to depression care among Latino men in a primary care setting: a qualitative study*. BMC Primary Care. 2024;25(1):30. doi:10.1186/s12875-024-02275-x

5. Wong EC, Torres VN, Martinez MO, Han B, Vue M, Derose KP. *A parish-based multilevel cluster randomized controlled trial to reduce stigma and mental health treatment disparities among Latino communities.* Contemporary Clinical Trials. 2023;125:107080. doi:10.1016/j.cct.2023.107080

6. Documet PI, Troyer MM, Macia L. *Social Support, Health, and Health Care Access Among Latino Immigrant Men in an Emerging Community*. Health Education & Behavior. 2019;46(1):137-145. doi:10.1177/1090198118763870

7. Ortega AN, McKenna RM, Pintor JK, et al. *Health Care Access and Physical and Behavioral Health Among Undocumented Latinos in California.* Medical Care. 2018;56(11):919. doi:10.1097/MLR.0000000000000985

8. Five Educational Learning Theories. Western Governors University. August 11, 2021. Accessed November 8, 2024. https://www.wgu.edu/blog/five-educational-learning-theories2005.html

9. Bulus M. *Difference between Two Means (Paired Samples t Test).* https://pwrss.shinyapps.io/index/

10. Matthew Ogrodnik, MS, CIP FKE PhD. *What is "Exempt" Human Subject Research, And What Does It Mean?* (2019 Edition). Accessed November 25, 2024. https://wwwapp.bumc.bu.edu/ocr/ClinicalResearchNewsletter/article.aspx?article=738

CURRICULUM VITAE