

2017-01

Has the phasing out of stavudine in accordance with changes in WHO guidelines led to a decrease in single-drug substitutions in first-line antiretroviral therapy for HIV in sub-Saharan Africa?

Brennan, A. T., Davies, M. A., Bor, J., Wandeler, G., Stinson, K., Wood, R., ... & Sikazwe, I. (2017). Has the phasing out of stavudine in accordance with changes in WHO guidelines led to a decrease in single-drug substitutions in first-line antiretroviral therapy for HIV in sub-Saharan Africa?. *AIDS (London, England)*, 31(1), 147-157. <https://doi.org/10.1097/QAD.0000000000001307>. <https://hdl.handle.net/2144/31120>

"Downloaded from OpenBU. Boston University's institutional repository."



Published in final edited form as:

AIDS. 2017 January 02; 31(1): 147–157. doi:10.1097/QAD.0000000000001307.

Has the phasing out of stavudine in accordance with changes in WHO guidelines led to a decrease in single-drug substitutions in first-line antiretroviral therapy for HIV in sub-Saharan Africa?

Alana T Brennan^{1,2,3,†}, Mary-Ann Davies⁴, Jacob Bor¹, Gilles Wandeler^{5,6}, Kathryn Stinson⁴, Robin Wood⁷, Hans Prozesky⁸, Frank Tanser⁹, Geoffrey Fatti¹⁰, Andrew Boulle⁴, Izukanji Sikazwe¹¹, Kara Wool-Kaloustian¹², Constantin Yiannoutsos^{12,13}, Valérie Leroy¹⁴, Nathalie de Rekeneire¹⁵, and Matthew P Fox^{1,2,3}

¹Department of Global Health, Boston University, Boston, MA, USA ²Health Economics and Epidemiology Research Office, Department of Internal Medicine, School of Clinical Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South ³Department of Epidemiology, Boston University School of Public Health, Boston, MA, USA ⁴Centre for Infectious Disease Epidemiology and Research, Cape Town, South Africa ⁵Department of Infectious Diseases, Bern University Hospital, University of Bern, Switzerland ⁶Institute of Social and Preventive Medicine, University of Bern, Switzerland ⁷The Desmond Tutu HIV Centre, Institute of Infectious Disease and Molecular Medicine, Faculty of Health Sciences, University of Cape Town, South Africa ⁸Division of Infectious Diseases, Department of Medicine, University of Stellenbosch & Tygerberg Academic Hospital, Cape Town, South Africa ⁹Africa Center for Health and Population Studies, University of Kwazulu-Natal, South Africa ¹⁰Kheth'Impilo AIDS Free Living, Cape Town, South Africa ¹¹Center for Infectious Disease Research in Zambia, Lusaka, Zambia ¹²Indiana University School of Medicine, Indianapolis, IN, USA ¹³Richard M. Fairbanks School of Public Health, Indiana University, Indianapolis, IN, USA ¹⁴INSERM U1027, Université Paul Sabatier Toulouse 3, Toulouse, France ¹⁵Université Bordeaux, ISPED, Centre INSERM U1219 Epidémiologie-Biostatistique, Bordeaux, France

Abstract

OBJECTIVE—We assessed the relationship between phasing out stavudine in first-line antiretroviral therapy (ART) in accordance with World Health Organization 2010 policy and single-drug substitutions (SDS) (substituting the nucleoside reverse transcriptase inhibitor (NRTI) in first-line ART) in sub-Saharan Africa.

DESIGN—Prospective cohort analysis (IeDEA-Multiregional) including ART-naïve, HIV-infected patients 16 years initiating ART between January 2005–December 2012. Before April 2010 (July 2007 in Zambia) national guidelines called for patients to initiate stavudine-based or zidovudine-based regimen, while after tenofovir or zidovudine replaced stavudine in first-line ART.

[†]Corresponding Author Details: Department of Global Health, Boston University, Crosstown Center, 3rd Floor, 801 Massachusetts Ave, Boston, MA 02118, Phone: 617-414-1479, abrennan@bu.edu.

METHODS—We evaluated the frequency of stavudine use and SDS by calendar year, 2004-2014. Competing risk regression was used to assess the association between NRTI use and SDS in the first 24 months on ART.

RESULTS—33,441 (8.9%; 95% CI: 8.7-8.9%) SDS occurred among 377,656 patients in the first 24 months on ART, close to 40% of which were amongst patients on stavudine. The decrease in SDS corresponded with the phasing out of stavudine. Competing risks regression models showed that patients on tenofovir were 20% to 95% less likely to require a SDS than patients on stavudine, while patients on zidovudine had a 75% to 85% decrease in the hazards of SDS when compared to stavudine.

CONCLUSION—The decline in SDS in the first 24 months on treatment appears to be associated with phasing out stavudine for zidovudine or tenofovir in first-line ART in our study. Further efforts to decrease the cost of tenofovir and zidovudine for use in this setting is warranted to substitute all patients still receiving stavudine.

Keywords

antiretroviral therapy; single-drug substitution; low- and middle-income countries; drug toxicities; drug side-effects

INTRODUCTION

In sub-Saharan Africa an estimated 11.4 million HIV-positive individuals were receiving antiretroviral therapy (ART) by mid-2015[1]. In low- and middle-income countries(LMICs), HIV treatment programs take a public health approach that utilizes twelve antiretrovirals in four drug classes[2,3]. Since treatment options are limited in LMICs and at least three drugs from two drug classes are typically needed for effectiveness, therapeutic options need to be maximized. One way to increase the life span of first-line ART in patients in LMICs is by decreasing rates of substituting individual drugs, typically for reasons of toxicity, within the regimen (“single-drug substitutions(SDS)”).

Prior to the World Health Organization(WHO) guidelines change in 2010, the most frequently used nucleoside reverse-transcriptase inhibitor(NRTI) alongside lamivudine was stavudine[2,3]. Stavudine is highly effective in treating HIV[4] but is associated with severe side-effects[5-7]. In 2009, as a result of stavudine’s poor side-effect profile, the WHO recommended replacing stavudine with tenofovir or zidovudine for initial HIV treatment[4]. In 2010, most governments in LMICs followed suit and began to phase out stavudine and replace it with tenofovir or zidovudine, for all new ART initiates[8]. Since there are fewer side-effects and drug toxicities associated with tenofovir and zidovudine than stavudine[9-12] the switch was expected to be accompanied by a marked decrease in SDS in first-line ART.

To date, several observational studies set in LMICs have compared rates of SDS amongst patients on stavudine-, tenofovir- and zidovudine-based regimens. All studies found that patients on tenofovir had about an 82% decrease in the risk of SDS(summary risk ratio (RR):0.18; 95% confidence interval (CI):0.15-0.20) compared to patients on stavudine[13-21], while patients on zidovudine, although at higher risk of SDS compared to

tenofovir, remained at lower risk of substitution compared to patients on stavudine(summary RR:0.41; 95%CI:0.37-0.45)[13-18].

While previous observational studies do suggest tenofovir and zidovudine are associated with fewer SDS compared to stavudine[13-21], many had important limitations. Two studies[13,14] had patient populations that initiated tenofovir prior to implementation of the 2010 WHO policy change. These patients would be more likely to have initiated tenofovir because of contraindications to stavudine or zidovudine leading to strong confounding by indication. We sought to use one of the largest HIV database in the world to assess whether or not the phasing out of stavudine in first-line ART in accordance with WHO 2010 policy decreased SDS in sub-Saharan Africa. This transition allows evaluation of the impact of a major policy change while accounting for secular trends in improvements in HIV treatment.

METHODS

Cohort Description

The International epidemiological Databases to Evaluate AIDS (IeDEA, www.iedea.org) is a world-wide National Institute of Health sponsored collaboration of HIV treatment cohorts. This study included cohorts from Southern Africa, East Africa and West Africa[22]. Data is collected on patients at the start of ART and at each follow-up visit. Clinic information includes demographic, clinical and HIV regimen data. Before April 2010(2007 in Zambia), if a patient experienced side-effects or toxicities related to stavudine or zidovudine, and was not in need of second-line therapy, the recommendation was to substitute stavudine with either zidovudine, if no related anemia or neutropenia was present, or abacavir and to substitute zidovudine with either stavudine or abacavir[23-29]. After April 2010(2007 in Zambia) patients initiated onto stavudine or zidovudine now had tenofovir, if no signs of renal-insufficiency were detected, while those initiated onto tenofovir could substitute with stavudine, zidovudine, or abacavir[23-29].

All IeDEA sites obtained ethical approval from relevant local institutions before contributing anonymized patient data. Approval for analysis of de-identified data was granted by Boston University's Institutional Review Board.

Study design

We performed a cohort analysis of data collected prospectively as part of routine care at clinics in the IeDEA multiregional collaboration. We included ART-naïve, HIV-infected patients 16 years old initiating first-line ART between January 1 2005-December 31 2012 for all countries except Nigeria, where patients initiating ART between January 1 2007-December 31 2012 were included as the roll out of ART started later. All patients had the potential for 24 months of follow-up. Prior to April 2010 national HIV treatment guidelines recommended the use of stavudine or zidovudine in first-line ART in all six included countries, thereafter, guidelines called for tenofovir or zidovudine to replace stavudine[23-27]. The only exception was in Zambia, which switched from stavudine or zidovudine to tenofovir in July 2007[28,29].

Study variables

All demographic (i.e. age, sex, clinic and country) and clinical (i.e. year of ART initiation, CD4 count, hemoglobin levels, weight, WHO stage, first-line NRTI (stavudine, zidovudine or tenofovir) and non-nucleoside reverse transcriptase inhibitor (NNRTI)(nevirapine or efavirenz) characteristics measured at ART initiation came from routinely collected clinic data. WHO staging was not available for East African countries.

The primary outcome variable was the proportion of patients who underwent a SDS in the first 24-months on ART. Follow-up time of 24-months was chosen as monitoring and time to development of toxicity/side-effects differ between drugs. Laboratory monitoring for tenofovir and zidovudine is conducted early on after treatment initiation, while for stavudine monitoring begins more often when the patient begins to develop clinical symptoms of toxicity(up to 24-48 months on ART[30]) diagnosed at a medical visit[2,3]. SDS was defined as substitution of the NRTI only within first-line ART. The reason for SDS were not available.

Statistical analysis

Patient characteristics at ART initiation were summarized with descriptive statistics and stratified by country. To look for trends in the use of stavudine in first-line ART and SDS over time, proportions of patients initiating stavudine or having a SDS in the first 24 months on ART were stratified by country and year of ART initiation and plotted from January 1 2005-December 31 2012, separately, with Nigeria being the exception, as data collection began in 2007. To test an additional hypothesis that tenofovir was being used among patients with contraindications to stavudine prior to guideline change, we looked at rates of SDS by NRTI over time.

Fine and Gray's competing risks regression method[31] was used to identify if the choice of NRTI in first-line ART was a predictor of SDS in the first 24 months on ART, accounting for attrition as competing risks and adjusted for age, sex, year of ART initiation, CD4 count, hemoglobin levels, WHO stage, and first-line NNRTI depending on country, with robust estimates at site level. We ran two models for each country. In both, we included all demographic and clinical characteristics at treatment initiation and year of ART initiation. The models differed as year of treatment initiation and NRTIs used are highly associated and therefore each model used only one of the two. Follow up time began at ART initiation and ended at the earliest of: 1) SDS; 2) initiation of second-line ART; 3) discontinuation of treatment 3) loss to follow-up (defined as not attending the clinic in the last 6 months); 4) death; 5) transfer; 6) completion of 24-months of follow-up; or 7) or date of dataset closure(December 31 2014).

We assessed interaction between gender and NRTI, CD4 count and NRTI, hemoglobin levels and NRTI, CD4 count and gender, WHO stage and gender and hemoglobin levels and gender on the additive scale by calculating the risk due to interdependence (R(I))[32].

Bayesian analysis

As this is not the first study on the topic and we can draw stronger conclusions when incorporating those prior results into our current analysis, we conducted a Bayesian analysis which allows incorporation of the result of prior knowledge about the relationship between the exposure and the outcome into the estimation of parameters. Our approach to Bayesian analysis[33] is essentially a weighted average incorporating the prior distribution (previous literature) and our data. To do this, point estimates and corresponding 95% confidence intervals (CI) for priors were obtained from previous publications assessing predictors of SDS[13-21]. Ratio measures for each potential predictor of SDS (age, sex, clinic, CD4 count, weight, hemoglobin levels, WHO stage, NRTI and NNRTI used in first-line regimen) were extracted from the existing literature. We performed a meta-analysis using random effects models, due to heterogeneity in estimates, to first create weighted summary estimates for each individual predictor, separately, from the existing literature (referred to as the “prior”). The same technique was used to calculate the summary estimates of each individual predictor, separately, from our data (referred to as the “likelihood”) and then combined the prior and likelihood estimates for each predictor, separately, to calculate a combined summary estimate (referred to as the “posterior”) and corresponding Bayesian credible intervals(CrI).

Sensitivity analysis

As we may have had unmeasured confounding in our population we conducted a multidimensional sensitivity analysis[34], by making assumptions about the strength of the effect of an unmeasured confounder on SDS and its prevalence in both patients on tenofovir or zidovudine (exposed) and those on stavudine (unexposed). We were interested in whether the confounder would overestimate the effect of exposure to tenofovir or zidovudine. We considered a confounder that would increase SDS and was more prevalent in patients on stavudine. We then back-calculated the relative risk we would have observed had we collected data on and adjusted for the purported confounder[34].

Multiple imputation

To account for missing data, we used multiple imputation by chained equations method using PROC MI in SAS[35] and assumed that the data were missing at random[36]. All models were fitted using 25 imputed datasets and estimated coefficients combined by averaging with the MIANALYZE procedure in SAS[37]. All clinical and demographic variables were included in the imputed models in addition to the outcome of SDS and indicator variables for death and loss to inform the missingness, but were not imputed. Appropriate standard errors were calculated using the within and between imputation standard errors of the estimates using Rubin’s rules[36]. Since models based on imputed results did not differ from the models on the original data we displayed the results based on the original data.

RESULTS

We included 377,656 patients in the analysis (Table 1), 24% initiated a stavudine-based ART regimen, ranging from from 25% in Zambia and Uganda to 60% in South Africa. Zambia

contributed the largest number of patients (n=205,140) and Nigeria the smallest (n=7,434). Demographic and clinical characteristics were similar across countries. Patients were predominately female (62.9%) with a median age of 35.2 years (interquartile range (IQR): 29.8-42.0) and a median time on treatment of 24.0 months (IQR:12.1-24.0), which did not differ by cohort. At ART initiation patients had a median CD4 count of 155 cells/mm³ (IQR: 74-241), with patients in South Africa having the lowest median (130 cells/mm³;IQR: 58-199) and Nigeria the highest (192 cells/mm³;IQR:91-312).

When stratified by year, gender, weight and age remained unchanged over time, while patients' cellular immunity at ART initiation improved in all countries. Additionally, the proportion of patients with tuberculosis and WHO III/IV stage declined over time in South Africa. Over 70% of patients in all countries remained alive and in care over 24 months on treatment. Overall attrition(combination of death and loss to follow-up) was 17.8% (95%CI: 17.7-17.9%) and fairly consistent across countries with, Zambia having the lowest rate of attrition in the first 24 months on ART at 15.3% (95%CI:15.2-15.5%) and Nigeria the highest at 23.5% (95%CI:22.5-24.4%)

Compliance with WHO guidelines: phasing out of stavudine in first-line ART

All countries, with the exception of South Africa where, in 2009, 95% of patients still initiated stavudine, began phasing out stavudine prior to the WHO guidelines making the change in 2010 (Figure 1). Zambia, Kenya and Uganda, began replacing stavudine with zidovudine in first-line ART as early as 2007 (2005 in Zambia), potentially in parallel with the WHO's recommendation for lower dose stavudine use (30mg instead of 40mg[38]). Tenofovir was introduced after the WHO recommended its use in first-line therapy in 2010 (2007 in Zambia), as such, within two years of the change in 2012, fewer than 10% of patients were being initiated on stavudine. Prior to 2010 in Cote d'Ivoire and Nigeria, stavudine and zidovudine were used interchangeably, while stavudine use decreased substantially, with less than 10% of patients initiating the drug after 2010, when both countries decided on zidovudine as the NRTI of choice in first-line ART. Tenofovir had yet to be introduced in first-line ART in Cote d'Ivoire or Nigeria before 2012 due to cost[3].

Decrease in SDS associated with the phase out of stavudine

While the WHO policy change was accompanied by a clear shift away from stavudine to tenofovir, the impact on single drug substitutions is less clear. Overall, SDS affected 8.8%(95%CI:8.7-8.9%) of patients in the first 24 months on ART with 38% of SDS related to stavudine compared to 49% and 13% related to zidovudine or tenofovir, respectively. The decrease in SDS was associated with the phasing out of stavudine in first-line ART, decreasing from an overall rate of 11.3%(95%CI:11.2-11.4%) prior to 2010 when 55% of patients were initiating stavudine to 5.4%(95%CI:5.3-5.5%) after 2010 when only 7.4% of patients initiated treatment with this drug. However, it is important to note that in all countries SDS began roughly two years prior to the WHO guideline change in 2010(Figure 2). For all countries competing risks regression models adjusted for year confirmed our results, showing a decrease in the hazards of SDS in accordance with the decrease in the use of stavudine(Supplementary Table 1).

SDS stratified by initiating NRTI and substitution NRTI

In addition to the variation observed in relation to the policy change, we also observed differences in rates of substitution by treatment regimen. Patients initiating stavudine(13.9%; 95%CI:13.7-14.1%) and zidovudine(12.0%; 95%CI:11.8-12.2%) had higher rates of single drug substitution compared to patients initiating tenofovir(2.8%; 95%CI:2.7-2.9%). Also, while rates of substitution decreased over time for patients on tenofovir, they remained stable for zidovudine patients and increased for those on stavudine(Supplementary Figure 1). All countries followed the national ART guidelines outlining the antiretrovirals eligible for substitution. These included tenofovir, stavudine, zidovudine or abacavir depending on NRTIs included in the first-line regimen with the occasional use of didanosine as an alternative(Supplementary Figure 2). Zidovudine was the most common SDS used for both tenofovir(50%; 95%CI:48.6-51.6%) and stavudine(67.9%; 95%CI:67.2-68.6%), while stavudine was the SDS for a zidovudine-based first-line regimen(65.1%; 95%CI: 64.2-66.0%).

Consistent with known toxicity patterns, SDS occurred earlier for patients on zidovudine(median 8.1 months after start of treatment; IQR:2.3-17.0) and tenofovir(median 10.2 months after start of treatment; IQR:3.6-18.1) compared to stavudine(median 14.2 months; IQR:7.3-19.6) after start of treatment.

Changing the NRTI used in first-line ART could explain the temporal trends in SDS

Adjusted competing risks regression models evaluating the association between choice of NRTI used in first-line ART and SDS helped us to confirm that the decrease in rates of SDS was associated with the phasing out of stavudine for tenofovir or zidovudine in first-line ART. Adjusted models showed that patients initiating tenofovir in Southern and East Africa were 20% to 95% less likely to undergo a substitution than patients initiating stavudine(Table 2). Posterior Bayesian estimates using an informative prior, showed close to an 80% reduction in the risk of SDS(posterior risk ratio (RR):0.21; 95% credible interval(CrI):0.20-0.22) when comparing tenofovir to stavudine (Table 3). With the exception of Zambia, where we saw an increase in the hazards of SDS when comparing zidovudine to stavudine(hazard ratio (HR):2.59; 95%CI:2.3-3.0) and Kenya, where we saw no association(HR:1.13; 95%CI:0.99-1.28)(Table 2), patients on zidovudine compared to stavudine in Uganda, Cote d'Ivoire and Nigeria had a 75% to 85% decrease in the hazards of SDS when compared to stavudine. Posterior Bayesian estimates using an informative prior, showed close to a 70% reduction in the risk of SDS(posterior RR:0.31; 95%CrI:0.30-0.33) (Table 3).

Our results also suggest that females compared to males have a 50% increase in the risk of SDS(posterior RR:1.48; 95%CrI:1.43-1.52)(Table 2 and Table 3). Since we believed changes in the trends of substitutions for sex could vary by NRTI used in first-line ART we calculated the risk due to interdependence(R(I)). With the exception of Zambia and Kenya, where the R(I) was essentially zero, in the other four countries showed a positive interdependence (Uganda 2%; Cote d'Ivoire 4%, South Africa 5% and Nigeria 6%). In other words, in Nigeria for example, 6% of SDS in women on stavudine is related to the dual action of female gender and stavudine. The risk of SDS in the doubly exposed(females

exposed to stavudine) was 41.9% vs the risk of 6.0% in the doubly unexposed (males unexposed to stavudine). We did not see any signs of interaction when assessing all other biological relationships using the R(I).

We saw inconsistent results when comparing nevirapine to efavirenz. In Zambia, Kenya, Uganda and Nigeria patients on nevirapine had a 10% to 70 % increase in the risk of SDS compared to those on efavirenz, while patients in South Africa and Cote D'Ivoire had a decrease in SDS of 20% and 40%, respectively. We also saw a decrease in the risk of SDS by 10% to 40% in all countries amongst patients with CD4 count <100 cells/mm³ compared to 100 cells/mm³.

Bias Analysis

Bias analyses simulating a confounder that would overestimate the effect of tenofovir or zidovudine vs. stavudine showed that in order for adjustment for an unmeasured confounder to bring our results close to null, the confounder would have to be present in 5% in those exposed to tenofovir or zidovudine, extremely common among patients exposed to stavudine (40%) and be a very strong predictor of SDS (RR >11.3), highly unlikely (Supplementary Table 2).

DISCUSSION

In the largest study to date, across multiple countries on the African continent, we show steady decrease in SDS corresponding to the phasing out of stavudine, in accordance with the WHO guidelines, from an overall rate of 11% prior to 2010 when 55% of patients were initiating stavudine to 5% after 2010 when only 7% of patients initiated treatment with this drug. Using Bayesian methods[33], although high heterogeneity between studies for the majority of estimates, we were able to estimate an 80% decrease in the risk of SDS when comparing tenofovir to stavudine, and 70% decrease when comparing stavudine to zidovudine, further highlighting the better safety profile associated with tenofovir and zidovudine compared to stavudine. Our results also showed a decrease in the rates of SDS from 2005 to 2012 for patients on tenofovir, providing evidence to support the notion that patients in earlier years where being initiated on tenofovir due to contraindications to stavudine or zidovudine and therefore at higher risk of toxicity/side-effects.

Time to substitution varied depending on the NRTI used in first-line ART, with zidovudine and tenofovir occurring early after treatment initiation and at a higher rate later on in follow-up amongst patients on stavudine. This is consistent with previous studies conducted in sub-Saharan Africa[13-21], however, it is important to note that time to substitution is partly a function of the frequency of monitoring, which differs for each NRTI. Laboratory monitoring for tenofovir and zidovudine is often conducted early on after treatment initiation, while for stavudine monitoring begins more often when the patient begins to develop clinical symptoms of toxicity diagnosed at a medical visit[2,3]. Additionally, although national guidelines for substitution of the NRTI and NNRTI within first-line ART were the same, the difference in estimates when comparing zidovudine to stavudine and also nevirapine to efavirenz could be due to variation in monitoring practices or the availability of NRTIs for substitution in each country.

Females, compared to males, had a 50% increase in the risk of SDS in our study, consistent with previous research[13,15-20]. By assessing effect measure modification on the additive scale, we also showed that depending on country, 2% to 6% , of SDS in women on stavudine was related to the dual action of female gender and stavudine. Although we could not confirm the reason for substitution in our study, previous research has reported women on stavudine are at higher risk of toxicity/side-effects than men[30,39,40] and that differences in risk of toxicity observed between genders could be related to differences in susceptibility or to a higher level of adherence to therapy achieved by women[30]. In addition to the size of our study population, we are the first to evaluate trends in SDS over almost a decade of treatment in public sector since the roll out of ART in 2004 in six countries in sub-Saharan Africa. However, our findings should be considered in light of the study limitations. First, this study represents patients from urban areas. Although some clinics are run out of tertiary hospitals, the majority operate at the primary care level, are led by nurses or clinical officers rather than physicians and are part of the public health care system of the country and may, therefore, not be generalizable to other clinics. Second, there was variability in the estimates of the association between NRTI used in first-line ART and SDSs by country. When comparing tenofovir to stavudine we saw a 20% to 95% decrease in the hazards of SDS and when comparing zidovudine to stavudine we saw an 85% reduction to a 200% increase in the hazards of SDS. Although national guidelines for substitution of the NRTI within first-line ART were the same in all six countries, the difference could be due to variation in monitoring practices at the level of the clinic or clinician or the availability of alternative NRTIs used for substitution. Third, due to the lack of documentation of reasons for SDS for almost 95% of events amongst patients with the event, we are likely underestimating the frequency and type of side effects due to less than perfect surveillance. There is a chance SDS in our study were driven by the policy change and not by side-effects/toxicity of stavudine. When conducted this study in our pediatric population we did see a substantial spike in SDS around the time of the guideline change as clinicians were substituting stavudine, regardless if the patient was tolerating stavudine well, with zidovudine or tenofovir. If substitutions were being driven by the guidelines in adults we believe we would see a similar increase, which is not present in these data. Fourth, patients with lower CD4 count (<100 cells/mm³) had upwards of 40% decrease in the risk of SDS over the follow-up period in some countries. As we do see higher rates of attrition in patients with a lower CD4 count (<100 cells/mm³ - 23.6% vs. 100 cells/mm³ - 14.2%), the decreased risk amongst patients with poorer immune status could be caused by survivor bias. Fifth, in order to strengthen the argument for further reduction in the cost in tenofovir and zidovudine for use in LMICs we recognize that it would be important to know the proportion of individuals who are symptomatic with toxicity related to stavudine-based therapy and did not experience a SDS. Unfortunately, conditions are poorly captured in the database preventing us from accurately assess symptoms amongst patients who did not have a SDS. However, Figure 1 shows that the policy change switching to tenofovir was the reason for the decline in SDSs in South Africa, as the country went from almost 100% of patients initiating stavudine prior to the 2010 guidelines changing to $<10\%$, which we believe allowed us to evaluate what a change in national policy to included tenofovir in first-line ART might do in other countries as they make the change. Sixth, it is possible that our populations differed with respect to some unmeasured confounder, as data on WHO staging were not available for Kenya and

Uganda. We may therefore have residual confounding in our estimates from those countries. However, our bias analysis suggests such an unmeasured confounder would be extremely unlikely in our cohort. Finally, multiple imputation helps make it possible to handle missing data routinely and improve the validity of research. However, deviations from the assumptions needed[35] could have led to unpredictable biases in our parameter estimates.

Conclusion

The decline in SDS in the first 24 months on treatment is associated with phasing out of stavudine for zidovudine or tenofovir in first-line ART in the countries in our study. When calculating the number needed to treat we found that to prevent one additional SDS event six patients would need to be treated over 24 months, which would be considered beneficial. Supporting further efforts to decrease the cost of tenofovir and zidovudine for use in LMICs.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Source of Support: Funding was provided by USAID under the terms of INROADS USAID-674-A-12-00029 and the National Institutes of Allergy and Infectious Diseases (NIAID) the IeDEA initiative Award Number 2U01AI069924, IeDEA-Southern Africa Award Number U01AI069924-10, IeDEA-East Africa Award Number U01AI0699-11, IeDEA-West Africa Award Number U01AI069919-10. This study is made possible by the generous support of the American people through the United States Agency for International Development (USAID) and the National Institutes of Health. The contents are the responsibility of the authors and do not necessarily reflect the views of USAID, NIH or the United States government.

References

1. Global Health Sector Response to HIV, 2000-2015: Focus on innovations in Africa. Available at: http://apps.who.int/iris/bitstream/10665/198065/1/9789241509824_eng.pdf
2. World Health Organization. Scaling up antiretroviral therapy in resource-limited settings: treatment guidelines for public health approach. 2003 Revisions.
3. World Health Organization. Antiretroviral therapy for HIV infection in adults and adolescents: recommendations for a public health approach - 2010 revision. 2010
4. Gallant JE, Staszewski S, Pozniak AL, DeJesus E, Suleiman JMAH, Miller MD, Coakley DF, Lu B, Toole JJ, Cheng AK. Efficacy and safety of Tenofovir DF vs stavudine in combination therapy in antiretroviral-naïve patients: a 3-year randomized trial. *JAMA*. 2004; 292:191–201. [PubMed: 15249568]
5. Domingos H, Cunha RVD, Paniago AMM, Martins DM, Elkhoury EB, Souza ASD. Metabolic effects associated to the highly active antiretroviral therapy (HAART) in AIDS patients. *Braz J Infect Dis*. 2009; 13:130–136. [PubMed: 20140358]
6. van Griensven J, Zachariah R, Rasschaert F, Mugabo J, Atté EF, Reid T. Stavudine- and nevirapine-related drug toxicity while on generic fixed-dose antiretroviral treatment: incidence, timing and risk factors in a three-year cohort in Kigali, Rwanda. *Trans R Soc Trop Med Hyg*. 2010; 104:148–153. [PubMed: 19732926]
7. Cournil A, Coudray M, Kouanfack C, Essomba CN, Tonfack CAD, Biwolé-Sida M, Delaporte E, Bork K, Laurent C. Reduced dose of stavudine and lipoatrophy in HIV-infected patients in Cameroon. *Antivir Ther*. 2010; 15:1039–1043. [PubMed: 21041920]
8. Global HIV/AIDS response. Epidemic update and health sector progress towards Universal Access. Progress Report 2011.

9. Gallant JE, Staszewski S, Pozniak AL, DeJesus E, Suleiman JMAH, Miller MD, Coakley DF, Lu B, Toole JJ, Cheng AK. Efficacy and safety of Tenofovir DF vs stavudine in combination therapy in antiretroviral-naïve patients: a 3-year randomized trial. *JAMA*. 2004; 292:191–201. [PubMed: 15249568]
10. Menezes CN, Crowther NJ, Duarte R, Van Amsterdam D, Evans D, Dickens C, Dix-Peek T, Rassool M, Prinsloo A, Raal F, Sanne I. A randomized clinical trial comparing metabolic parameters after 48 weeks of standard- and low-dose stavudine therapy and tenofovir disoproxil fumarate therapy in HIV-infected South African patients. *HIV Med*. 2014 Jan; 15(1):3–12. Epub 2013 Aug 28. DOI: 10.1111/hiv.12074 [PubMed: 23980620]
11. Arribas JR, Pozniak AL, Gallant JE, DeJesus E, Gazzard B, Campo RE, et al. Tenofovir disoproxil fumarate, emtricitabine, and efavirenz compared with zidovudine/lamivudine and efavirenz in treatment-naïve patients: 144-week analysis. *J Acquir Immune Defic Syndr*. 2008; 47(1):74–8. [PubMed: 17971715]
12. Haubrich, Richard H., Riddler, Sharon A., DiRienzo, A Gregory, Komarow, Lauren, Powderly, William G., Klingman, Karin, Garren, Kevin W., Butcher, David L., Rooney, James F., Haas, David W., Mellors, John W., Havlir, Diane V. Metabolic Outcomes in a Randomized Trial of Nucleoside, Nonnucleoside and Protease Inhibitor-Sparing Regimens for Initial HIV Treatment AIDS. Author manuscript; available in PMC. 2010 Jun 1.
13. Velen K, Lewis JJ, Charalambous S, et al. Comparison of tenofovir, Zidovudine, or Stavudine as part of first-line antiretroviral therapy in a resource-limited-setting: a cohort study. *PLoS One*. 2013 May 14.8(5):e64459. [PubMed: 23691224]
14. Bygrave H, Ford N, Van CG, et al. Implementing a tenofovir-based first-line regimen in rural Lesotho: clinical outcomes and toxicities after two years. *J Acquir Immune Defic Syndr*. 2011 Mar 1; 56(3):e75–8. [PubMed: 21164354]
15. Chi BH, Mwango A, Giganti M, et al. Early clinical and programmatic outcomes with tenofovir-based antiretroviral therapy in Zambia. *J Acquir Immune Defic Syndr*. 2010; 54:63–70. [PubMed: 20009765]
16. Brennan AT, Maskew M, Ive P, et al. Increases in regimen durability associated with the introduction of tenofovir at a large public-sector clinic in Johannesburg, South Africa. *J Int AIDS Soc*. 2013 Nov 19.16:18794. [PubMed: 24256692]
17. Njuguna C, Orrell C, Kaplan R, et al. Rates of switching antiretroviral drugs in a primary care service in South Africa before and after introduction of tenofovir. *PLoS One*. 2013 May 22.8(5):e63596. [PubMed: 23717451]
18. Inzaule S, Otieno J, Kalyango J, Nafisa L, Kabugo C, Nalusiba J, Kwaro D, Zeh C, Karamagi C. Incidence and predictors of first line antiretroviral regimen modification in western Kenya. *PLoS One*. 2014 Apr 2.9(4):e93106. [PubMed: 24695108]
19. Labhardt ND, Sello M, Lejone T, Ehmer J, Mokhantso M, Lynen L, Pfeiffer K. Adoption of new HIV treatment guidelines and drug substitutions within first-line as a measure of quality of care in rural Lesotho: health centers and hospitals compared. *Trop Med Int Health*. 2012 Oct; 17(10): 1245–54. [PubMed: 22845835]
20. Boule A, Orrel C, Kaplan R, Van Cutsem G, McNally M, Hilderbrand K, Myer L, Egger M, Coetzee D, Maartens G, Wood R. Substitutions due to antiretroviral toxicity or contraindication in the first 3 years of antiretroviral therapy in a large South African cohort. *International Epidemiological Databases to Evaluate Aids in Southern Africa Collaboration. Antivir Ther*. 2007; 12(5):753–60. [PubMed: 17713158]
21. Takuva S, Louwagie G, Zuma K, Okello V. Durability of First line Antiretroviral Therapy: Reasons and Predictive Factors for Modifications in a Swaziland Cohort. *J Antivir Antiretrovir*. 2012; 4:014–020.
22. Egger M, Ekouevi DK, Williams C, et al. Cohort Profile: the international epidemiological databases to evaluate AIDS (IeDEA) in sub-Saharan Africa. *Int J Epidemiol*. 2012 Oct; 41(5): 1256–64. [PubMed: 21593078]
23. National Department of Health, Republic of South Africa. The South African Antiretroviral Treatment Guidelines. 2010. Available at: <http://www.doh.gov.za/docs/factsheets/guidelines/art.pdf>

24. Ministry of Health, Government of Kenya. Guidelines for antiretroviral drug therapy in Kenya. 42011. Available at: <http://healthservices.uonbi.ac.ke/sites/default/files/centraladmin/healthservices/Kenya%20Treatment%20Guidelines%202011.pdf>
25. Ministry of Health, Government of Uganda. National Antiretroviral treatment guidelines for Adults, adolescents and children. Third Edition. MOH 2009; 2009. Available at: <http://apps.who.int/medicinedocs/documents/s19249en/s19249en.pdf>
26. Federal Ministry of Health. Government of Cote D'Ivoire. Guidelines for antiretroviral drug therapy in Cote D'Ivoire. 2010
27. Federal Ministry of Health. Department of Public Health Nigeria. National guidelines for HIV and AIDS treatment and care in adolescents and adults. 2010. Available at http://www.who.int/hiv/pub/guidelines/nigeria_art.pdf
28. Ministry of Health, Republic of Zambia. Adult and Adolescent Antiretroviral Therapy Protocols. 2010. Available at: http://www.who.int/hiv/pub/guidelines/zambia_art.pdf
29. Mwinga, A. Inclusion of a tenofovir-based first-line regimen in Zambia: a bold step forward?; Paper presented at: 2007 HIV/AIDS Implementer's Meeting; Kigali, Rwanda. 2007;
30. Pujades-Rodríguez, Mar, Dantony, Emmanuelle, Pinoges, Loretxu, Ecochard, René, Etard, Jean-François, Carrillo-Casas, Esther, Szumilin, Elisabeth. for the AIDS Working Group of Médecins Sans Frontières. Toxicity Associated with Stavudine Dose Reduction from 40 to 30 mg in First-Line Antiretroviral Therapy. PLoS One. 2011; 6(11):e28112. [PubMed: 22132226]
31. Fine, Jason P., Gray, Robert J. A Proportional Hazards Model for the Sub distribution of a Competing Risk. JSTOR. :496–509.
32. Rothman KJ. Interactions between causes. Mod Epidemiol. 1986:311–326.
33. MacLehose, Richard F., Hamra, Ghassan B. Applications of Bayesian methods to epidemiologic research. Current Epidemiology Reports 1.3. 2014:103–109.
34. Lash, TL., Fox, MP., Fink, AK. Applying Quantitative Bias Analysis to Epidemiologic Data. Springer; 2009.
35. The MI Procedure. Available: <http://support.sas.com/rnd/app/papers/miv802.pdf>
36. Rubin, DB. Multiple imputation for Nonresponse in Surveys. New York: John Wiley & Sons; 1987.
37. The MIANALYZE Procedure. Available: <http://support.sas.com/rnd/app/papers/mianalyzev802.pdf>
38. World Health Organization. Addendum to 2006 WHO Guidelines on Antiretroviral Therapy for HIV Infection in Adults and Adolescents. Geneva: 2007.
39. Emery J, Pick N, Mills EJ, Cooper CL. Gender differences in clinical, immunological, and virological outcomes in highly active antiretroviral-treated HIV-HCV coinfecting patients. Patient Prefer Adherence. 2010 May 13.4:97–103. [PubMed: 20517470]
40. Kempf MC, Pisu M, Dumcheva A, Westfall AO, Kilby JM, Saag MS. Gender differences in discontinuation of antiretroviral treatment regimens. J Acquir Immune Defic Syndr. 2009 Nov 1; 52(3):336–41. [PubMed: 19654551]




Figure 1. Trends in the stavudine use in first-line antiretroviral therapy for treatment naïve patients stratified by country and year since ART guidelines changed (N=377,656)
 *Dashed lines represents the change in national guidelines to introduce zidovudine or tenofovir into first-line ART (2007 for Zambia and 2010 for all other countries)




Figure 2. Proportion of single-drug substitutions over the first 24-months on antiretroviral therapy stratified by year of since ART guidelines changed for all countries (N=377,656)
 *Dashed lines represents the change in national guidelines to introduce zidovudine or tenofovir into first-line ART (2007 for Zambia and 2010 for all other countries)

Table 1

Clinical and demographic characteristics at initiation of antiretroviral therapy and outcomes over 24-months of follow-up amongst adults initiated on treatment stratified by country (N=377,656), 2005-2012

Characteristics	Country						Total (N=377656) n (%)
	South Africa (n=47290) n (%)	Zambia (n=205140) n (%)	Kenya (n=73547) n (%)	Uganda (n=31645) n (%)	Cote d'Ivoire (n=12600) n (%)	Nigeria (n=7434) n (%)	
Nucleoside Reverse Transcriptase Inhibitor							
stavudine	28698 (60.7)	51289 (25.0)	36399 (49.5)	7846 (24.8)	7011 (55.6)	2775 (37.3)	87456 (23.2)
tenofovir	18592 (39.3)	115108 (56.1)	17686 (24.1)	4796 (15.2)	-	-	156182 (41.4)
zidovudine	-	38743 (18.9)	19462 (26.5)	19003 (60.1)	5589 (44.4)	4659 (62.7)	134018 (35.5)
Non-nucleoside Reverse Transcriptase Inhibitor at First-line ART Initiation							
efavirenz	38563 (81.6)	88873 (43.3)	22835 (31.1)	10486 (33.1)	5129 (40.7)	717 (9.6)	166603 (44.1)
nevirapine	8727 (18.5)	116267 (56.7)	50712 (69.0)	21159 (66.9)	7471 (59.3)	6717 (90.4)	211053 (55.9)
Gender							
male	17660 (37.3)	78965 (38.5)	25304 (34.4)	11598 (36.7)	4337 (34.4)	2354 (31.7)	140218 (37.1)
female	29630 (62.7)	126175 (61.5)	48243 (65.6)	20047 (63.4)	8263 (65.6)	5080 (68.3)	237438 (62.9)
Age (years)							
18-24.9	2842 (6.0)	18905 (9.2)	6868 (9.3)	3094 (9.8)	637 (5.1)	483 (6.5)	32829 (8.7)
25-29.9	8213 (17.4)	36398 (17.7)	11257 (15.3)	5708 (18.0)	1724 (13.7)	1423 (17.1)	64723 (17.1)
30-39.9	21005 (44.4)	89589 (43.7)	28534 (38.8)	13529 (42.8)	5406 (42.9)	3017 (40.6)	161080 (42.7)
40-49.9	10948 (23.2)	42571 (20.8)	17943 (24.4)	6940 (21.9)	3475 (27.6)	1644 (22.1)	83521 (22.1)
50	4282 (9.1)	17677 (8.6)	8945 (12.2)	2374 (7.5)	1358 (10.8)	867 (11.7)	35503 (9.4)
median (IQR)	35.0 (30.0-42.0)	34.9 (29.5-41.4)	36.2 (30.1-43.8)	34.9 (29.3-41.1)	37.2 (31.5-43.7)	35.4 (29.9-43.0)	35.2 (29.8-42.0)
CD4 count (cells/mm³)							
0-49	8734 (18.5)	10533 (5.1)	12205 (16.6)	5286 (16.7)	1923 (15.3)	850 (11.4)	151617 (40.2)
50-99	7082 (15.0)	12070 (5.9)	9720 (13.2)	3743 (11.8)	1290 (10.2)	733 (9.9)	39531 (10.5)
100-199	14097 (29.8)	25615 (12.5)	19523 (26.5)	8558 (27.0)	2756 (21.9)	1457 (19.6)	34638 (9.2)
200	9955 (21.1)	30204 (14.7)	23609 (32.1)	9396 (29.7)	3859 (30.63)	2841 (38.2)	72006 (19.1)
missing	7422 (15.7)	126718 (61.8)	8490 (11.5)	4662 (14.7)	2772 (22.0)	1553 (20.9)	79864 (21.2)
median (IQR)	130 (58-199)	165 (87-255)	155 (70-252)	156 (69-230)	163 (70-253)	192 (91-312)	155 (74-241)
Hemoglobin (ug/dL)							

Characteristics	Country							Total (N=377656) n (%)
	South Africa (n=47290) n (%)	Zambia (n=205140) n (%)	Kenya (n=73547) n (%)	Uganda (n=31645) n (%)	Cote d'Ivoire (n=12600) n (%)	Nigeria (n=7434) n (%)		
10	19930 (42.1)	19300 (9.4)	36382 (49.5)	14363 (45.4)	4189 (33.3)	2954 (39.7)	89975 (23.8)	
<10	6152 (13.0)	9117 (4.4)	15689 (21.3)	3337 (10.6)	3954 (31.4)	1372 (18.5)	34295 (9.1)	
missing	21208 (44.9)	176723 (86.2)	21476 (29.2)	13945 (44.1)	4457 (35.4)	3108 (41.8)	233352 (65.3)	
median (IQR)	11.6 (10.0-13.0)	10.9 (9.4-12.2)	11.2 (9.6-12.8)	11.9 (10.5-13.3)	10.0 (7.8-11.0)	10.0 (9.0-12.0)	10.0 (8.0-11.4)	
Weight (kg)	median (IQR)	54.1 (48.0-61.0)	55.0 (49.0-62.0)	54.0 (48.0-61.0)	57.3 (66.0-50.0)	61.0 (53.0-70.0)	55.0 (49.0-62.4)	
World Health Organization Stage								
I/II	21965 (46.5)	95394 (46.5)	-	-	1136 (9.0)	4605 (62.0)	170757 (45.2)	
III/IV	24980 (52.8)	96787 (47.2)	-	-	3316 (26.3)	1503 (20.2)	172351 (45.6)	
missing	345 (0.7)	12959 (6.3)	-	-	8148 (64.7)	1326 (17.8)	12959 (6.3)	
Time (months)	median (IQR)	24.0 (12.9-24.0)	24.0 (10.0-24.0)	24.0 (12.8-24.0)	24.0 (11.8-24.0)	24.0 (8.7-24.0)	24.0 (12.1-24.0)	
Vital status over 24-months of follow-up								
Death	n (%)	12611 (6.2)	5121 (7.0)	1674 (5.3)	660 (5.2)	119 (1.6)	24113 (6.4)	
Loss	n (%)	5863 (12.4)	18859 (9.2)	10768 (14.6)	3761 (11.9)	1626 (21.9)	1626 (21.9)	
Attrition	n (%)	9791 (20.7)	31470 (15.3)	15889 (21.6)	5435 (17.2)	2968 (23.6)	67298 (17.8)	
Transfers	n (%)	3757 (7.9)	31271 (15.2)	2563 (3.5)	2618 (8.3)	285 (2.3)	40524 (10.7)	
Alive	n (%)	33742 (71.4)	142399 (69.4)	55095 (74.9)	23592 (74.6)	9347 (74.2)	269834 (71.4)	
Primary outcome over 24-months of follow-up								
Single-drug substitution	n (%)	5686 (12.0)	13574 (6.6)	8074 (11.0)	2654 (8.4)	1512 (12.0)	1695 (22.8)	33441 (8.85)

Table 2 Competing risks regression models examining predictors of single-drug substitution over 24-months on ART in all countries (N=377,656)

Variable measured at ART initiation	Single-Drug Substitution					
	Southern Africa		East Africa		West Africa	
	SOUTH AFRICA Adjust Hazards Ratio (95% CI)	ZAMBIA Adjust Hazards Ratio (95% CI)	KENYA Adjust Hazards Ratio (95% CI)	UGANDA Adjust Hazards Ratio (95% CI)	COTE D'IVOIRE Adjust Hazards Ratio (95% CI)	NIGERIA Adjust Hazards Ratio (95% CI)
NRTI in first-line ART regimen						
stavudine	Reference	Reference	Reference	Reference	Reference	Reference
tenofovir	0.12 (0.10-0.16)	0.80 (0.49-1.29)	0.47 (0.44-0.50)	0.05 (0.02-0.11)	-	-
zidovudine	-	2.61 (1.83-3.71)	1.13 (1.00-1.28)	0.13 (0.08-0.19)	0.24 (0.17-0.48)	0.14 (0.12-0.16)
NNRTI in first-line ART regimen						
efavirenz	Reference	Reference	Reference	Reference	Reference	Reference
nevirapine	0.79 (0.64-0.96)	1.30 (1.03-1.66)	1.22 (0.97-1.55)	1.03 (0.67-1.59)	0.58 (0.48-0.69)	1.92 (1.45-2.55)
Sex						
Male	Reference	Reference	Reference	Reference	Reference	Reference
Female	1.84 (1.59-2.15)	1.36 (1.22-1.53)	1.31 (1.25-1.38)	1.11 (0.93-1.34)	1.51 (1.15-1.97)	1.12 (0.95-1.31)
Age (years)						
16-24.9	0.63 (0.62-0.64)	0.84 (0.70-1.01)	0.85 (0.74-0.97)	1.04 (0.74-1.44)	0.62 (0.50-0.77)	1.05 (0.80-1.40)
25-29.9	0.84 (0.76-0.92)	0.88 (0.80-0.97)	0.85 (0.73-0.99)	0.95 (0.90-1.01)	0.88 (0.69-1.12)	0.84 (0.68-1.04)
30-39.9	Reference	Reference	Reference	Reference	Reference	Reference
40-49.9	1.26 (1.17-1.38)	1.16 (1.04-1.29)	1.26 (1.15-1.37)	1.05 (0.98-1.14)	1.22 (1.13-1.32)	0.90 (0.75-1.08)
50	1.44 (1.38-1.51)	1.12 (0.93-1.36)	1.56 (1.40-1.73)	0.97 (0.87-1.09)	1.23 (1.00-1.51)	0.93 (0.74-1.17)
CD4 count (cells/mm³)						
200	Reference	Reference	Reference	Reference	Reference	Reference
100-199	0.95 (0.93-0.97)	0.94 (0.84-1.07)	0.92 (0.83-1.01)	0.85 (0.63-1.14)	0.80 (0.67-0.96)	1.00 (0.84-1.20)
50-99	0.96 (0.88-1.05)	0.88 (0.77-1.00)	0.88 (0.76-1.00)	0.70 (0.49-1.01)	0.87 (0.63-1.19)	0.93 (0.74-1.17)
<50	0.81 (0.79-0.84)	0.79 (0.65-0.97)	0.80 (0.71-0.92)	0.59 (0.42-0.85)	0.73 (0.59-0.90)	0.72 (0.57-0.91)
Weight (kg)						
10	1.01 (1.02-1.03)	1.00 (0.99-1.01)	1.01 (1.00-1.02)	1.00 (0.99-1.01)	1.02 (1.00-1.03)	1.00 (0.99-1.01)
Hemoglobin (ug/dL)						
10	Reference	Reference	Reference	Reference	Reference	Reference

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

		Single-Drug Substitution					
		Southern Africa		East Africa		West Africa	
Variable measured at ART initiation		SOUTH AFRICA Adjust Hazards Ratio (95% CI)	ZAMBIA Adjust Hazards Ratio (95% CI)	KENYA Adjust Hazards Ratio (95% CI)	UGANDA Adjust Hazards Ratio (95% CI)	COTE D'IVOIRE Adjust Hazards Ratio (95% CI)	NIGERIA Adjust Hazards Ratio (95% CI)
<10		0.88 (0.77-1.00)	1.96 (1.69-2.26)	1.06 (1.03-1.10)	1.04 (0.76-1.42)	1.09 (0.86-1.38)	0.75 (0.63-0.89)
WHO stage							
I/II	Reference	Reference	Reference	Reference	Reference	Reference	Reference
III/IV	1.11 (1.05-1.16)	1.10 (0.97-1.26)	-	-	1.20 (0.79-1.81)	1.06 (0.90-1.25)	

Table 3
 Bayesian analysis: prior, likelihood and posterior estimates for predictors of single-drug substitution

Variable	Bayesian Estimates					
	Prior Risk Ratio (95% CI)	Prior I ² , p-value	Likelihood Risk Ratio (95% CI)	Likelihood I ² , p-value	Posterior Risk Ratio (95% CrI [*])	Posterior I ² , p-value
NRTI						
stavudine	Reference		Reference		Reference	
tenofovir	0.18 (0.15-0.20)	93.1%, p=0.000	0.22 (0.20-0.23)	99.3%, p=0.000	0.21 (0.20-0.22)	86.2%, p=0.007
zidovudine	0.41 (0.37-0.45)	97.1%, p=0.000	0.21 (0.18-0.25)	98.8%, p=0.000	0.31 (0.30-0.33)	96.4%, p=0.000
NNRTI						
efavirenz	Reference		Reference		Reference	
nevirapine	0.87 (0.74-1.01)	0.0%, p=0.521	0.86 (0.78-0.95)	89.3%, p=0.000	0.86 (0.79-0.93)	0.0%, p=0.902
Sex						
Male	Reference		Reference		Reference	
Female	1.76 (1.69-1.83)	96.9%, p=0.000	1.33 (1.28-1.38)	90.7%, p=0.000	1.48 (1.43-1.52)	99.0%, p=0.000
Age (years)						
16-24.9	0.96 (0.80-1.12)	0.0%, p=0.797	0.83 (0.77-0.90)	73.2%, p=0.002	0.85 (0.79-0.91)	54.1%, p=0.140
25-29.9	0.96 (0.80-1.12)	0.0%, p=0.797	0.91 (0.87-0.95)	9.5%, p=0.355	0.91 (0.87-0.95)	0.0%, p=0.552
30-39.9	Reference		Reference		Reference	
40-49.9	0.91 (0.75, 1.07)	0.0%, p=0.971	1.00 (0.97-1.02)	93.8%, p=0.000	1.00 (0.97-1.02)	15.7%, p=0.276
50	0.92 (0.76-1.08)	0.0%, p=0.741	1.07 (1.01-1.12)	94.1%, p=0.000	1.05 (1.00-1.11)	66.9%, p=0.082
CD4 count (cells/mm³)						
200	Reference		Reference		Reference	
100-199	0.93 (0.82-1.03)	0.0%, p=0.633	0.93 (0.88-0.98)	0.0%, p=0.721	0.93 (0.88-0.98)	0.0%, p=1.000
50-99	0.93 (0.82-1.05)	0.0%, p=0.688	0.87 (0.81-0.94)	0.0%, p=0.462	0.88 (0.83-0.94)	0.0%, p=0.373
<50	0.97 (0.85-1.09)	31.6%, p=0.223	0.78 (0.72-0.83)	0.0%, p=0.462	0.81 (0.76-0.86)	87.4%, p=0.005
Weight (kg) **	1.00 (0.50-1.50)	-	1.01 (1.00-1.01)	71.6%, p=0.014	1.01 (1.00-1.01)	0.0%, p=0.969
Hemoglobin (ug/dL)						
10	Reference		Reference		Reference	
<10	0.98 (0.96-1.01)	51.3%, p=0.152	1.05 (1.02-1.08)	95.6%, p=0.000	1.01 (0.99-1.03)	91.9%, p=0.000
WHO stage						

Bayesian Estimates

Variable	Prior Risk Ratio (95% CI)	Prior I ² , p-value	Likelihood Risk Ratio (95% CI)	Likelihood I ² , p-value	Posterior Risk Ratio (95% CrI*)	Posterior I ² , p-value
I/II	Reference 1.08 (1.0-1.16)	27.0%, p=0.241	Reference 1.04 (0.95-1.15)	94.6%, p=0.000	Reference 1.06 (1.00, 1.13)	0.0%, p=0.540
III/IV						

* CrI, Credible Interval (Bayesian)

** Only posterior estimate that was estimated with a flat prior (RR=1.0; 95% CI: 0.5-1.5)